

Trauma, Violence, & Abuse

<http://tva.sagepub.com/>

Child Sexual Abuse, Links to Later Sexual Exploitation/High-Risk Sexual Behavior, and Prevention/Treatment Programs

Kevin Lalor and Rosaleen McElvaney

Trauma Violence Abuse 2010 11: 159 originally published online 2 August 2010

DOI: 10.1177/1524838010378299

The online version of this article can be found at:

<http://tva.sagepub.com/content/11/4/159>

Published by:



<http://www.sagepublications.com>

Additional services and information for *Trauma, Violence, & Abuse* can be found at:

Email Alerts: <http://tva.sagepub.com/cgi/alerts>

Subscriptions: <http://tva.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

Citations: <http://tva.sagepub.com/content/11/4/159.refs.html>

Child Sexual Abuse, Links to Later Sexual Exploitation/High-Risk Sexual Behavior, and Prevention/Treatment Programs

Kevin Lalor¹ and Rosaleen McElvaney¹

Abstract

This paper reviews the literature on the nature and incidence of child sexual abuse, explores the link between child sexual abuse and later sexual exploitation, and reviews the literature on prevention strategies and effective interventions in child sexual abuse services. Our understanding of the international epidemiology of child sexual abuse is considerably greater than it was just 10 years ago, and studies from around the world are examined. Childhood sexual abuse can involve a wide number of psychological sequelae, including low self-esteem, anxiety, and depression. Numerous studies have noted that child sexual abuse victims are vulnerable to later sexual revictimization, as well as the link between child sexual abuse and later engagement in high-risk sexual behaviour. Survivors of child sexual abuse are more likely to have multiple sex partners, become pregnant as teenagers, and experience sexual assault as adults. Various models which attempt to account for this inter-relationship are presented; most invoke mediating variables such as low self-esteem, drug/alcohol use, PTSD and distorted sexual development. Prevention strategies for child sexual abuse are examined including media campaigns, school-based prevention programmes, and therapy with abusers. The results of a number of meta-analyses are examined. However, researchers have identified significant methodological limitations in the extant research literature that impede the making of recommendations for implementing existing therapeutic programmes unreservedly.

Keywords

etiology, sexual abuse, treatment/intervention

Unfortunately, evidence worldwide indicates that children and adolescents are at risk from sexual predators, exploiters, and opportunists. Incidence varies from time to time and from place to place, depending on individual, familial, and societal circumstances, but the sexual exploitation of young people appears to be a universal phenomenon. The extant research also shows that family members, relatives, neighbors, or those known and trusted by the child, typically perpetrate abuse.

Scientific literature on the nature and incidence of child sexual abuse (CSA) has existed for almost 150 years. Masson (1992) details the recognition of CSA by physicians in 19th-century France. For example, he describes Tardieu's *A Medico-Legal Study of Assaults on Decency*, which noted that 9,125 people were accused of rape or attempted rape of children between 1858 and 1869.

In more recent times, widespread recognition of child abuse is typically dated to the United States in the early 1960s. Initial concerns were with the "battered child syndrome." This term was first used by pediatrician C. Henry Kempe at a 1961 symposium of the American Pediatric Society (Kempe, Silverman, Steele, Droegmueller, & Silver, 1962). The identification of this syndrome came about through observations of the large numbers of unexplained broken bones and bruises in children on pediatric

wards. The growing awareness and concern regarding, first, child physical abuse and, later, CSA in the medical and child protection literature in the United States in the 1960s and 1970s has been well documented elsewhere (DeMause, 1991; Schultz, 1982).

The purpose of this article is to review the literature on the nature and incidence of CSA internationally, with a particular focus on perpetrators; to explore the link between CSA and (later) exploitation outside the family; to review the literature on CSA prevention strategies; and to review the literature on effective interventions and best practices in CSA services.

Overview of the Nature and Incidence of CSA

In an early international perspective on child abuse, the Foreword to Korbin's (1981) *Child abuse and neglect*:

¹ Department of Social Sciences, Dublin Institute of Technology

Corresponding Author:

Kevin Lalor, Department of Social Sciences, Dublin Institute of Technology, 40-45 Mountjoy Square, Dublin, Ireland.
 Email: kevin.lalor@dit.ie

Cross-cultural perspectives states “child abuse . . . has become a serious social problem in the United States and in some other industrialized countries, yet it occurs infrequently or not at all in many of the world’s societies” (p. vii). We now know that this is not the case.

In an important study, Finkelhor (1994) aimed to illustrate the international epidemiology of CSA and found that research studies in 19 countries produced findings similar to North American research (incidence rates ranging from 7% to 36% for women and 3% to 29% for men), thus undermining the idea that CSA is “peculiarly American.” He concluded:

Studies from a variety of countries suggest that sexual abuse [of children] is indeed an international problem. In every locale where it has been sought, researchers have demonstrated its existence at levels high enough to be detected through surveys of a few hundred adults in the general population . . . As such epidemiological findings are available for more and more countries, the responsibility of proof shifts to anyone who would argue that sexual abuse is rare or nonexistent in their locale. (1994, p. 412)

Finkelhor (1994) noted the preponderance of results from Caucasian, Western, Christian countries and the relative lack of data from middle-Eastern, African, or Far-Eastern countries. Lachman (1996) also noted the absence of research on child abuse in Africa¹ where it is “overshadowed by political and economic problems, lack of resources, the enormity of the phenomenon, and the lack of a research culture and research experience” (1996, p. 543). However, he did note a growing number of publications from Africa in international journals.

Today, although many gaps remain, we have better information about the nature and incidence of CSA throughout the world. In the material below, selected studies are summarized. Data on incidence and perpetrators, in particular, are highlighted. First, however, it is important to highlight some methodological considerations.

CSA Research: Definitions and Methodological Considerations

The methodological limitations of child abuse research have long been recognized (Briere, 1992; Runyan, 2000; Widom, 1988). In addition to the inherent secrecy and sensitivity of the issue, numerous methodological issues must be noted.

First, researchers use a wide variety of definitions of “childhood sexual abuse.” Many general population surveys define it as “unwanted sexual contact” without asking for specific details of the behavior. Some studies distinguish between “contact abuse” and “noncontact abuse” or “penetrative abuse” and “non-penetrative abuse.” Studies use different age cutoff points (before the age of 15, 16, 17, or 18) to define CSA. Studies that focus on more broadly based definitions of CSA not surprisingly indicate higher prevalence rates.

Second, studies have been conducted among a range of different samples; college samples, convenience samples, clinical

samples, and national probability samples. The latter are the least commonly studied, given the cost involved but yield the most generalizable data.

Third, and perhaps most importantly, prevalence rates are accurate only to the extent that respondents are willing to discuss the topic with researchers or complete questionnaires. Consider the response rates reported in the following studies: McGee, Garavan, de Barra, Byrne, and Conroy (2002); Ireland, national sample of adults—71%; Chen, Dunne, and Han (2004); China, sample of high school students—70.5%; Briere and Elliott (2003); United States, national sample of adults—64.8%; Steel and Herlitz (2005); Sweden, national sample of adults—59%.

As we can see, nonresponse rates of 30% are common, thus the findings from the literature are based on the 70% of people who typically agree to participate.

It is important that we are cognizant of the above issues when interpreting the findings of CSA studies. As noted by Pereda, Guilera, Forns, and Gomez-Benito (2009), following their review of 39 prevalence studies across 28 countries and 5 continents, difficulties remain in reaching a consensus on the definition of child maltreatment and sexual abuse (Manly, 2005, cited in Pereda et al., 2009). For the purpose of this review, studies that refer to CSA, with or without behaviorally specific definitions, will be included. Where authors have clarified their definition of CSA, this will be referred to wherever possible. The objective of this review is to be inclusive, acknowledging the limitations of this approach.

CSA Prevalence Worldwide

As noted above, there is a growing body of well-designed studies from around the world published in international peer-reviewed journals. An exhaustive review is beyond the scope of this article. Instead, a geographically diverse selection of studies carried out in recent years will be briefly presented in Table 1 below.

As we can see, rates for penetrative CSA are higher for girls than for boys. For female children/adolescents, rates for penetrative abuse range from 0.3% (China); between 1.6% and 3% (South Africa); 4% (Israel); 5.6% (Ireland); 11% (Tanzania); to a high of 18% reporting “unwanted sexual intercourse” in one Ethiopian study.

Data on perpetrators are not always reported, but the studies above show that (for *all* types of CSA) “strangers” make up only a minority of perpetrators. For example, 1% (New Zealand); 2% (United Kingdom); 10% (Swaziland); 16% (Israel); 21% (strangers and recent acquaintances; South Africa); and 36% (Ethiopia).

Accordingly, very high proportions of perpetrators were well known to the child; 86% were family members in the New Zealand study described above; 76% of perpetrators were family members or known to the child (Ireland); 78% were boyfriends, neighbors, or a male relative (Swaziland); and, in South Africa, 64% were teachers, relatives, or boyfriends.

Table 1. "Headline" Findings on Child Sexual Abuse (CSA)

Country	Study	Incidence	Perpetrators
UK	May-Chahal and Cawson (2005), <i>n</i> = 2,869, 18–24 years old	10% (6% male; 15% female) reported a form of contact CSA before age 16 2% (1% male; 3% female) reported sexual intercourse against their wishes before age 16	1% of the total sample reported contact CSA by parent/carer; 2% reported contact CSA by relative; 8% reported contact CSA by other known person; and 2% reported contact CSA by stranger/person just met Not reported
France	King, Guilbert, Ward, Arwidson, and Noubary (2006), <i>n</i> = 12,256 adults	1.3% (0.7% male; 2.1% female) a forced sexual relationship ("touching" or "attempted rape" or "rape") before age 18 Females: 20.4% contact sexual abuse before age 17, 5.6% penetrative abuse before age 17 Males: 16.2% contact sexual abuse before age 17, 2.7% penetrative abuse before age 17	Against females: 24% family members; 52% known to victim; 24% strangers Against males: 14% family members; 66% known to victim; 20% strangers Not reported
China	Chen et al. (2004), <i>n</i> = 2,300 high school students	Females: 16.7%, "unwanted sexual experience" before age 16 (6.5% touching or fondling; 0.3% unwanted sexual intercourse) Males: 10.5%, "unwanted sexual experience" before age 16 (2.8% touching or fondling; 0.1% unwanted sexual intercourse)	Not reported
Bedouin-Arabs in Southern Israel	Elbedour, Abu-Bader, Onwuegbuzie, Abu-Rabia, and El-Aassam (2006), <i>n</i> = 217 female adolescents	31% some form of CSA between the ages of 14 and 18 13% fondling; 4% penetrative abuse	Limited data provided. 16% strangers
New Zealand	Fanslow et al. (2007), <i>n</i> = 2,855 females	23.5% in the urban (Auckland) area and 28.2% of women in the rural (Waikato) region "sexually touched or made to do something sexual they did not want to" before age 15 32.3% of females and 14.2% of males, at least one episode of childhood sexual abuse (kissed or touched in a sexual way) before 18. 52.8% of child abuse incidents involved penetrative abuse	86% male family members 1% described as "strangers"
USA	Briere and Elliott (2003), <i>n</i> = national sample of 935 adults	33.3%, some form of sexual violence before age 18 9.1% "coerced intercourse" before age 18 4.9% "forced intercourse" before age 18	46.8% "incest" Against females: 93% by at least one male; 9% by at least one female Against males: 39% by at least one female, 70% by at least one male
Swaziland	CDCP and UNICEF (2007), <i>n</i> = 1,920 females aged 13–24 years	68.7%, some form of sexual violence before age 18 18.1% "coerced intercourse" before age 18 4.9% "forced intercourse" before age 18	35.6% husband or boyfriend, 27.1% a man/boy from the victim's neighborhood, 15.7% a male relative other than a father, stepfather or husband, 10.1% a stranger
Ethiopia	Woriku et al. (2006), <i>n</i> = 323 female secondary school students (aged 12 to 20 years)	68.7%, some form of sexual abuse "in their lifetime" 18.1% unwanted sexual intercourse	36.7% school mates, 16.7% family members (comprising 1.4% father, 9.3% step father, 2.3% elder brother, 0.9% uncle, 5% other family members) and 15.8% neighbors Limited data provided
Tanzania	McCann, Lalor and Katabaro (2006), <i>n</i> = 487 university students	27% of the sample (31% of females and 25% of males) one or more unwanted sexual experiences before the age of 18. Females: unwanted fondling, 27.9%; sexual intercourse, 11.2%; oral sex, 5.6% Males: unwanted fondling, 13.4%; sexual intercourse, 8.8%; oral sex, 6.2%	Limited data provided
South Africa	Jewkes, Levin, Mbananga, and Bradshaw (2002), <i>n</i> = 11,735 women aged 15–49 years	1.6% forced or persuaded to have sex against their will before age 15. 3% of 15- to 19-year-olds, forced or persuaded to have sex against their will before age 15	33% school teachers, 21% relations, 21% strangers or recent acquaintances, 10% boyfriends

Table 2. Incidence and Perpetrators of Childhood Sexual Abuse Against Females

Location	Percentage Reporting SA Before Age 15 ^a	Family ^b	Acquaintance ^c	Stranger ^d	Not Identified
Bangladesh city	7.4	10.9	14.3	69.7	5.9
Bangladesh province	1	*	*	*	*
Brazil city	11.6	66.3	14.1	8.7	14.1
Brazil province	8.7	54.1	30.6	12.9	11.8
Ethiopia province	7	*	*	*	*
Japan city	13.8	9.2	20.6	69.5	5.3
Namibia city	21.3	46.6	34.2	16.4	4.1
Peru city	19.5	53.6	16.7	25	19.2
Peru province	18.1	41.4	20	22.8	24.1
Samoa	1.8	23.3	33.3	33.3	16.7
Serbia and Montenegro city	4.2	28.6	25	39.3	10.7
Thailand city	8.9	7.7	12	58.1	24.8
Thailand province	4.9	21.7	11.7	30	38.3
United Republic of Tanzania city	12.2	27.8	29.1	13.9	32.9
United Republic of Tanzania province	9.5	18.3	41.7	18.3	28.3

Source: García-Moreno et al. (2005).

Note. ^aThe researchers asked questions related to child sexual abuse in face-to-face interviews, and by asking them to anonymously mark a card. In some regions, this produced quite different results. The results presented here are the researchers' "best estimate" of true incidence. ^b"Family" includes father, step-father, male, and female family members. ^cTeacher, friend of family, boyfriend, someone at work, religious leader. ^dIncludes police/soldier.

*Percentage based on less than 20 respondents suppressed.

These findings should not be seen as comparisons in prevalence across countries due to differences in study design, definitions, and sampling.

World Health Organization (WHO) Multi-Country Study

In addition to the individual studies mentioned above, a recent multi-country study is important to note. García-Moreno, Jansen, Ellsberg, Heise, and Watts (2005) report on a WHO multi-country study on women's health and domestic violence, including experiences of childhood sexual abuse. The report contains data from more than 24,000 survey respondents in 10 countries. This represents one of the largest scale studies of women's experiences of CSA (and other forms of violence) worldwide.

Results regarding CSA (defined as someone touching them sexually, or making them do something sexual they did not want to, before the age of 15) and the identity of perpetrators are shown in Table 2 below.

There are a number of important patterns to note here. First, the percentage of respondents reporting CSA varies widely worldwide, from a high of 21.3% in Namibia "city," to a low of 1% in Bangladesh "province." More research is needed to determine the risk and protection factors at work here.

It is also interesting to note the variance between these figures and some of the other studies we have looked at earlier. For example, Table 2 above shows that 7% of the sample in Ethiopia "province" reported "being touched sexually or being made to do something sexual they did not want to do" before age 15 years. We saw earlier that Worku, Gebremariam, and Jayalakshmi (2006) reported that a sample of 12- to 20-year-old Ethiopian schoolgirls reported a very high proportion (18%) experiencing unwanted sexual intercourse "in your lifetime." Clearly, the age cutoff of 15 in the WHO study will show lower

prevalence rates relative to other studies on CSA, which typically use a higher cutoff age (typically, before age 18).

Also of note is the identity of perpetrators. It is difficult to see any pattern here. In some regions, family members and acquaintances accounted for the bulk of the abuse (80.4% in Brazil "province;" 84.7% in Brazil "city;" and 80.8% in Namibia). By comparison, in Thailand "city," family and acquaintances only accounted for 19.7% of perpetrators.

In some regions, "strangers" are significant abusers (69.7% in Bangladesh "city" and 69.5% in Japan "city"). By contrast, they account for only 8.7% of abusers in Brazil "city."

Prevalence rates noted in this table are comparable to those cited in Finkelhor (1994) and a follow-up review conducted by Pereda et al. (2009). In both reviews, the majority of studies (67% and 50%, respectively) report prevalence rates for sexual abuse of male children of below 10% whereas in the case of females, most studies (38 and 34%, respectively) locate the prevalence between 10% and 20%. Another 29% of studies in both reviews report a prevalence rate of CSA among women of between 30% and 40%. Many of the studies reviewed defined CSA as either contact or noncontact sexual experiences between a person younger than 18 years of age and an adult or other person at least 5 years older or sexual experiences resulting from coercion, regardless of the age of the other person.

What Is the Link Between CSA and Later Sexual Exploitation?

The purpose of this section is to describe the sequelae of CSA. First, general mental health sequelae are briefly described. The remainder of this section focuses on the link between CSA and later sexual revictimization. The link between CSA and high-risk sexual behavior is also considered.

The Negative Sequelae of CSA

CSA includes a broad range of behaviors, which can be perpetrated across a broad range of intrafamilial and extrafamilial relationships, and there is considerable variability in the duration and frequency of the abuse (Paine & Hansen, 2002). This multifaceted nature accounts for the heterogeneity of children's reactions to abuse, ranging from severe psychological impact to no evidence of negative psychological sequelae (Kendall-Tackett, Williams, & Finkelhor, 1993).

Citing various reviews (e.g., Briere & Runtz, 1993; Finkelhor, 1990; Polusny & Follette, 1995), Briere and Elliott (2003) summarize two decades of research which:

suggest with relative unanimity that childhood physical and sexual abuse have a wide number of psychological sequelae. Among these are low self-esteem, anxiety, depression, anger and aggression, posttraumatic stress, dissociation, substance abuse, sexual difficulties, somatic preoccupation and disorder, self-injurious or self-destructive behavior, and most of the various symptoms and behaviours seen in those diagnosed with borderline personality disorder. (p. 1207)

For example, Dube et al. (2005) surveyed 17,337 adult members of a health plan in California regarding childhood experiences and current health and social problems. Overall, 16% of male respondents and 25% of females reported some type of contact sexual abuse before age 18. The results showed a markedly greater likelihood of social/health problems in adulthood among CSA victims, for both women and men. Victims were several times more likely to have alcohol problems, use illicit drugs, attempt suicide, and have current marriage/family problems.

Severity of CSA (penetrative vs. non-penetrative) is associated with elevated risk of social or health problems in adulthood. The authors conclude "exposure to CSA among both genders is common, and acts as a strong risk factor for multiple types of mental health, behavioral, and social outcomes similarly for adult men and women" (Dube et al., 2005, p. 434).

It is important to note that a significant proportion of children do not experience psychopathology following sexual abuse. Collishaw et al. (2007) report on data from the longitudinal Isle of Wight study in the United Kingdom. Approximately 10% of respondents (now in middle age, 44–45 years) reported having experienced physical or sexual abuse in childhood while about 8% reported severe or repeated physical or sexual abuse. Of those reporting abuse, a substantial percentage (about 45%) reported no psychopathology over their adult lives. The authors attribute this to factors such as parental care, peer relationships in adolescence, personality, and the quality of adult relationships.

Link Between CSA and Sexual Revictimization

In addition to socioemotional and mental health sequelae, numerous studies have highlighted that CSA victims are

vulnerable to later sexual revictimization. Most studies do not distinguish between intra- and extrafamilial CSA. Rather, they seek to investigate effects in adulthood, or correlates, of any form of childhood sexual, physical, or emotional maltreatment.

An early study was conducted by Russell (1986) who found that 63% of a sample of women who experienced intrafamilial sexual abuse before age 14, also experienced rape or attempted rape after age 14 (compared to 35% of those who did not report childhood sexual abuse), suggesting that childhood sexual abuse doubled the risk of sexual revictimization. Research over the past decade has consistently replicated Russell's findings with rates of victimization varying from study to study, ranging from 13.7% (Humphrey & White, 2000) to 27% (McGee et al., 2002). Classen, Palesh, and Aggarwal (2005) reviewed approximately 90 empirical studies and concluded that CSA is a significant risk factor for sexual revictimization. A review by Messman-Moore and Long (2003) found that CSA survivors are between 2 and 11 times more likely to experience adult assault compared to non-victims. Steel and Herlitz' (2005) study in Sweden of 2,810 adults found that those men and women who had experienced child or adolescent sexual abuse were at higher risk for adult sexual assault (8% vs. 1%).

Roodman and Clum (2001) reviewed 19 empirical studies of females, where at least one incidence of childhood sexual abuse was followed by a subsequent incident of adult sexual victimization. They found an effect size for revictimization of .59 and that the effect size varied from very low to very high, reflecting the variety of methods and definitions used. For example, those studies with an older average sample age showed a larger effect size (not surprisingly, as participants have had more time in which to be revictimized). The definitions of child and adult abuse were also found to be important. The effect size average for studies using broad definitions of child abuse was .38. This increased to .64 for those studies using a "contact" sexual abuse definition. McGee et al.'s (2002) high rate of 27% was based on penetrative CSA and penetrative sexual violence in adulthood. However, the risk of contact non-penetrative violence was 24% as compared with respective risks of 10% (penetrative) and 3% (contact non-penetrative) for those with no experience of abuse in childhood.

Sexual revictimization in adolescence appears to significantly increase the risk of sexual assault in adulthood. Humphrey and White (2000) report on a 5-year longitudinal study of sexual assault among a sample of 1,569 female college students. They found that victims of childhood assault were significantly more likely to experience either moderate (unwanted or verbal coercion) or severe (attempted or completed rape) adolescent victimization than non-victims. Respondents who had an experience of rape or attempted rape in their adolescent years were 13.7 times more likely to experience rape or attempted rape in their first year of college.

Classen et al. (2005) point out that more severe forms of CSA, and the coexistence of physical abuse, increase the risks for later revictimization. They also found that sexually abused children were more likely to be abused as adolescents; who were in turn more likely to be sexually

revictimized as adults. Widom, Czaja, and Dutton (2008) report the findings of a prospective study of children physically and sexually abused between 1967 and 1971, and a comparison group of non-abused children matched for sex, age, race, and socioeconomic status. Both groups were followed up and interviewed between 2000 and 2002 (at a mean age of 39.5 years). Those who experienced abuse/neglect in childhood had significantly higher rates of sexual assault/abuse in adulthood (47.7% vs. 28.6%).

Some studies report whether the revictimization in adulthood is perpetrated by a partner or a non-partner, influencing risk rates. A study of New Zealand of 2,855 women by Fanslow, Robinson, Crengle, and Perese (2007) found that women who reported childhood sexual abuse were 3½ times more likely to report sexual violence by a non-partner in their lifetime (23.2% vs. 6.4%). In the sample of Dunkle et al. (2004) of 1,395 women attending antenatal clinics in South Africa, 74.8% reported physical/sexual partner violence and 15.6% reported adult sexual assault by a non-partner. The equivalent prevalence for participants who did not report child sexual assault or forced first intercourse was 52.2% and 5.3%, respectively.

As is the case with prevalence rates of CSA, differences have also been found between men and women in revictimization risk rates. McGee et al. (2002) found that women had an 11% and 12% higher risk of penetrative and contact non-penetrative sexual violence, respectively, than men in their study.

The research is not unequivocal in establishing the link between sexual victimization in childhood and later sexual assault. Mandoki and Burkhart (1989) examined the relationship between child and adult sexual victimization among a sample of 282 female undergraduates (mean age 19.5 years). Unlike much of the research in this area, they found that those sexually victimized as children were *not* more likely than non-victims to be sexually assaulted as adults.

Link Between CSA and High-Risk Sexual Behavior

In addition to the link between CSA and revictimization, numerous studies have examined the relationship between CSA and later engagement in high-risk sexual behavior (such as having multiple sexual partners), prostitution, and sexual promiscuity.

Fergusson, Horwood, and Lynskey (1997) studied a birth cohort of 520 females in New Zealand from birth to age 18. They found clear correlations between CSA and high-risk sexual behaviors. For example, 58.6% of those who had experienced penetrative CSA had more than 5 sexual partners before age 18, compared to 13.3% who reported no form of CSA. Zierler et al. (1991) found that men and women who had been raped or forced to have sex as a child or teenager were twice as likely to have had multiple partners than those who were not assaulted. Other high-risk sexual behaviors found to be significantly more common in their abused sample included changing partners frequently and engaging in sex with casual

acquaintances. Krahé et al. (1999, cited in Messman-Moore & Long, 2003) found that the number of partners predicted sexual victimization, with women who reported CSA having had a significantly higher number of partners (average of 5.6 vs. 2.9 reported by women without histories of CSA), which was in turn related to higher rates of later victimization. Steel and Herlitz (2005) found that victims of child or adolescent sexual abuse were at higher risk of engagement in group sex compared to a sample of adults who had not experienced such abuse (12% vs. 6%).

Unprotected sexual intercourse has also been found to be significantly more prevalent in young women reporting CSA (Fergusson et al., 1997). Sexually transmitted diseases are an obvious concern when considering multiple sex partners and unprotected sex. Wingood and DiClemente (1997) found that in their sample of 165 female African Americans in a socioeconomically deprived area of San Francisco, 13.3% had experienced forced sex before the age of 16. This group was 1.4 times more likely to report having a sexually transmitted disease and 2.4 times more likely to report having more than 2 lifetime sexually transmitted diseases.

Early onset of consensual sexual activity has been associated with a history of CSA in women (Fergusson et al., 1997; Noll, Trickett, & Putnam, 2003). In the study by Noll et al., the average age of first voluntary intercourse in their female sample was 14 years, compared to 15.5 years in their matched comparison group. Early age at first intercourse has also been found to be associated with a history of child and adolescent abuse in Sweden (Steel & Herlitz, 2005) and China (Chen et al., 2004). The study of 2,810 adults by Steel and Herlitz found that 61% of child and adolescent sexual abuse (CASA) victims reported first intercourse prior to 17 years, compared to 42% of non-CASA victims. Related to this, teenage pregnancy has also been found to be more prevalent in studies of women (Fergusson et al., 1997). Zierler et al. (1991) found that women in their study were 2½ times more likely to have become pregnant as a teenager than those who were not sexually assaulted as a child or teenager.

Prostitution has been found to be more prevalent in studies of both women and men who experienced penetrative abuse in childhood or adolescence (Steel & Herlitz, 2005; Zierler et al., 1991). In the study by Zierler et al., women were four times more likely to report having worked as a prostitute. Widom and Kuhns (1996) found childhood sexual abuse to be a significant predictor of prostitution in female subjects. Overall, 10.73% of their abuse/neglect group had engaged in prostitution, compared to 5.6% of the control group. For females, the difference between the abuse/neglect group and the control group were 8.93% versus 2.87%, and for males it was 12.54% versus 8.03%. Potter, Martin, and Romans (1999) compared a group of 29 prostitutes, with a sample of similarly aged females from the general population. The former were significantly more likely than the latter to have experienced penetrative sexual abuse before age 16 (25% vs. 4.7%).

Bagley and Young (1987) found that 73% of former prostitutes had experienced CSA compared to 29% of a control group

from the general population. The former were significantly more likely to report penetrative sexual abuse before age 16 (25% vs. 4.7%). Furthermore, they found the sexual abuse experienced by the prostitute group was more frequent and serious in nature, involved more serious assault, more often involved different individuals, began at a significantly earlier age and continued for longer. They also found that 71% of the former prostitutes had “devastated self-esteem” compared to 7% of the control group.

It cannot be said, however, that CSA *causes* later high-risk behaviors and revictimization. There may be other mediating factors (such as parental conflict and parental psychopathology) that are associated with sexual abuse experiences and high-risk behavior and/or victimization.

Both the study by Widom and Kuhns (1996) and the study by Noll et al. (2003) were prospective studies from early childhood to early adulthood over a 10-year period, using matched comparison groups. The former study involved a sample size of 82 in the abused group and 84 in the matched control group while the latter study had a sample of 1,196 subject (676 abused and/or neglected and 520 control subjects). Noll et al. (2003) also used an explicit definition of CSA, a well-matched comparison group and they succeeded in retaining over 95% of the sample. A prospective design allows the effects of sexual abuse to be “disentangled” from other potential sources of trauma, such as poverty and chaotic family background.

Mediating Factors in the Link Between Childhood Sexual Abuse and Revictimization and High-Risk Sexual Behaviors

Respondents in the study by Fergusson et al. reporting CSA were significantly more likely to come from disturbed family/home environments, characterized by instability, physical punishment, poor parental attachment, and use of illicit drugs/alcohol. Similarly, Potter et al. (1999) found significant differences between their prostitute group and their control group regarding parental separation (38% of the prostitute group, compared to 8.5% of the control group). However, the association between CSA and variables such as multiple partners, unprotected intercourse, consensual intercourse before age 16, and rape/attempted rape remained statistically significant even allowing for these family variables. Fergusson et al. (1997) conclude that such findings suggest a possible cause and effect link in which CSA leads to increased sexual risk taking in adolescence and to early onset of sexual intercourse.

Widom and Kuhns (1996) speculate that running away and being alone on the streets and thus being vulnerable to pimps and pornographers may contribute to the links found between childhood sexual abuse and later high-risk sexual behaviors. Potter et al. (1999) compared the family backgrounds of 29 prostitutes, with a sample of similarly aged females from the general population. The former were significantly more likely than the latter to have experienced penetrative sexual abuse before age 16 (25% vs. 4.7%). The sex worker group were

considerably more likely to leave home early (55% had left home by age 17, compared to 26% in the general population).

Widom and Kuhns note the lack of attention in the research to separating out the effects directly related to sexual abuse from those possibly due to preexisting difficulties for the child, the family, or the stress associated with disclosure so that the unique contribution of childhood abuse and neglect can be realized.

Fergusson and Mullen (cited in Fleming, Mullen, Sibthorpe, & Bammer, 1999) note that the influence of a positive family environment operates at two levels. First in facilitating the development of the child in terms of resilience and second in providing a nurturing and supportive environment in which the child can recover from the impact of the abuse and reestablish the security needed for normal development.

Noll et al. (2003) suggest that stigmas associated with childhood sexual abuse “may result in an inability to glean nonsexual or emotional rewards from relationships, therefore placing victims in potentially risky or exploitative sexual situations” (p. 582).

Stoltz et al. (2007) found a significant relationship between child maltreatment (sexual, physical, and emotional) and later involvement in prostitution among a sample of 361 drug-using street-involved youth in Vancouver, Canada. They suggest two explanations for this relationship. First, those children who are sexually victimized “develop psychologically and emotionally in ways that make them vulnerable to continuing sexual predation.” Second, they speculate that “childhood sexual abuse may create a propensity in the victim toward risk-taking behaviors (substance abuse, running away from home) . . . that in turn lead to situations in which survival sex work is one of few remaining options” (p. 1218).

Stoltz and colleagues offer similar explanations for the susceptibility of emotionally abused youth to become involved in trading sex, thus emphasizing that CSA in and of itself does not “cause” later involvement in prostitution. Rather, it is a component of a generally traumatic, negative developmental experience.

Nevertheless, numerous studies report that juvenile prostitutes identify the sexual abuse they experienced as a child as a major factor in their decision to become involved in prostitution. For example, 70% of the subjects in the study by Silbert and Pines (1982) in the United States and 62% of those in a study by Bagley and Young (1987) in Canada reported that childhood sexual abuse affected their decision to become involved in prostitution. However, there is no evidence that any but a small proportion of children who are sexually victimized subsequently become involved in trading sex for money or goods. In other words, while Flowers (1998) can maintain that “nearly all girl prostitutes have been sexually molested, assaulted, or physically abused before entering the profession” (p. 82), the reverse does not hold. In Sweden, Steel and Herlitz (2005) found that 1% of the sample of 268 CASA victims reported having engaged in “sex work,” compared to 0% of the sample of 2,625 non-victims. This does not suggest a strong predisposition to involvement in prostitution among CASA victims.

In one African study, Slonim-Nevo and Mukuka (2007) surveyed 3,360 10- to 19-year-olds in Zambia regarding their experiences of physical and sexual abuse by family members and AIDS-related knowledge, attitudes, and behavior. They found that abuse by family members was a significant predictor of engagement in high-risk behaviors. The higher the level of sexual and physical abuse in the family, the higher the probability was of engaging in high-risk behavior. These findings were independent of various sociodemographic factors. They argue that intrafamilial abuse should not be ignored not only because of the negative impacts on the development of children but because it also reduces the likelihood of them being able to protect themselves from HIV/AIDS. The cultural context in sub-Saharan Africa where violence/coercion in early sexual relationships and exchange sex is so prominent may mask the relationship between early abusive sexual experiences and later high-risk behaviors such as multiple partners, exchange sex, prostitution, and repeat sexual assaults and rape (Lalor, 2008).

What are the Reasons for CSA Victims' Vulnerability to Later Revictimization and High-Risk Sexual Behavior?

The studies above do not always speculate as to *why* CSA victims are vulnerable to revictimization and risky sexual behavior. As many studies into CSA and its effects are cross-sectional in design, it is not possible to identify causal factors. Numerous mediating variables have been proposed, such as low self-esteem, alcohol use, dissociation, low danger awareness, and social isolation.

Possible Explanations for the Revictimization of CSA Victims. Cloitre's sociodevelopmental approach (1998, cited in Arata, 2002) proposes that abuse in childhood interferes with the development of affect regulation and interpersonal relatedness, which in turn affects women's awareness of danger and ability to respond to threatening situations. The model of Gold, Sinclair, and Balge (1999, cited in Arata, 2002) suggests that CSA leads to impairments in attachment, psychological symptoms, and negative attributes and coping behaviors, resulting in behaviors that increase the risk of rape.

Polusny and Follette (1995) proposed a conceptual model based on the theory of "emotional avoidance." From this perspective, many of the long-term correlates of CSA are the individual's efforts to minimize or reduce negative emotional states such as guilt, shame, fear and rage, and associated thoughts and memories. Over time, emotional avoidance behaviors (such as dissociation, substance abuse, and self-mutilation) "are negatively reinforced by the reduction or suppression of the intense affective responses associated with sexual abuse experiences" (p. 158). In this context, high-risk sexual behaviors (such as frequent, indiscriminate, and compulsive sexual behaviors) serve to modulate emotional pain by providing more pleasurable, distress-incompatible input (Briere & Runtz, 1993, cited in Polusny & Follette, 1995). This emotional avoidance may lead to revictimization. For example, chronic dissociative coping

behaviors may lead survivors to ignore or minimize social cues in dangerous adult situations (Walker et al., 1992, cited in Polusny & Follette, 1995). Sandberg, Matorin, and Lynn (1999, cited in Messman-Moore & Long, 2003) note the high levels of posttraumatic stress disorder (PTSD) among CSA survivors and suggest this can interfere with information processing of danger cues.

Steel and Herlitz (2005) hypothesize a possible pathway from CASA to sexual risk behavior, with psychological symptoms/disorders as a mediator. Psychological sequelae to CASA, such as depressive symptoms, poor self-esteem, lack of assertiveness, poor self-worth, and PTSD, may lead to future risk behavior due to "(1) feelings of unworthiness; (2) the inability to be assertive and prevent unwanted sexual advances; (3) avoidance or inability to comprehend emotionally laden information concerning sexuality; or (4) having competing needs for affection and acceptance" (p. 1150). They conclude, "future research is warranted to both understand the mechanisms underlying the relationship between CASA and SRBs and to explore other childhood and adolescent experiences that may predict SRBs as an adolescent or adult" (p. 1151).

Slonim-Nevo and Mukuka (2007) also speculate as to the mediating variables that link intrafamilial CSA and engagement in HIV-risk sexual behavior, they note:

Depression and other psychological problems may lead adolescents to a sense of hopelessness and futility about living. Such a perspective may thus lead to a sense of indifference about sickness and even death. Consequently, these adolescents engage in immediately satisfying sexual activities and fail to protect their future." (p. 156)

There is some empirical support for the role of dissociation as a mediating variable. Meadows, Jaycox, Orsillo, and Foa (1997, cited in Messman-Moore & Long, 2003) suggest that withdrawing from potentially harmful situations may be due to difficulties in risk recognition and acting on potential risk. They found no difference between previously victimized and non-victimized women (physical or sexual assault at any age) for risk recognition. However, the previously assaulted group reported that they would leave a hypothetical scenario (or "scene") significantly later than individuals with no history of assault.

Scenes with an acquaintance or someone known to the subject (rather than a stranger) were "left" significantly later by subjects with a history of assault. Furthermore, subjects without a history of assault also endorsed leaving the scene prior to physical contact, in contrast to previously abused subjects who usually did not endorse leaving until after physical contact had been made (Meadows et al., 1997, cited in Messman-Moore & Long, 2003, p. 550).

Many authors emphasize the need for prospective studies to enquire into the causal pathways from childhood sexual abuse to revictimization (Sanders & Moore, 1999; Arata, 2000, 2002). Messman-Moore and Long (2003) argue for a shift of

Table 3. Factors Hypothesized to be Related to Sexual Revictimization

Ecological Level	Factor (or variable)
Ontogenic development	The initial victimization experience (possible effects include traumatic sexualization, alcohol and drug abuse, dissociative disorders, low self-esteem, powerlessness, stigmatization, learned expectancy for victimization, social isolation, running away from home, deviance, and early/premarital pregnancy). Early family experiences ^a (possible correlates include family breakdown, disorganization and dysfunction, marital dysfunction, unsupportive parents, and patriarchal structure)
Microsystem	Exposure risk (factors increasing this risk include traumatic sexualisation, dissociative disorder, alcohol abuse, involvement with deviant activities, stigmatization, and low self esteem) Increased risk of perpetrator acting aggressively due to perception of victim as easy target, feeling justified in behaving aggressively, and victim's decreased ability to respond assertively and effectively to unwanted sexual advances
Exosystem	Lack of resources (related factors include low socioeconomic status, unsafe living conditions, early childbearing, single motherhood, and divorce). Lack of alternatives (due to weak family ties or support and social isolation)
Macrosystem	Cultural tendency to blame victim. Good girl/bad girl construction of femininity

^a As noted by Messman-Moore and Long (2003), the original ecological model proposed by Bronfenbrenner conceived the family environment as part of the microsystem.

focus from the victim to interpersonal and cultural factors within society:

“Our theories of revictimization will shape what we will find empirically. If we focus on the victim, we will find within her and her behavior explanations for revictimization . . . We will never fully understand revictimization while the microscope remains focused solely on the victim.” (p. 566)

Grauerholz (2000) proposes an ecological model to explain child sex abuse victims' increased risk of being sexually victimized later in life. The focus is on female victims as “research on male sexual revictimization is nonexistent” (p. 5). She notes that explanations frequently focus on individualistic factors, facilitating a “victim-blaming” interpretation; she argues that a comprehensive explanation must incorporate interpersonal and sociocultural factors. After Bronfenbrenner (1979) and Belsky (1980), an ecological perspective is proposed, which incorporates ontogenic development (one's individual personal history), the microsystem (e.g., family relations), the exosystem (larger social systems in which the individual or family is located), and the macrosystem (cultural norms and institutions). Thus, a rich contextual understanding is gained of the individual, interpersonal, community, and societal forces causing repeat sexual victimization. Grauerholz (2000) proposes a range of factors hypothesized to be mediating variables between childhood sexual abuse and later sexual exploitation (revictimization; Table 3).

Grauerholz (2000) notes the existence of a number of models or theoretical frameworks at the individual level that seek to explain how childhood sexual abuse affects a person's personality and behavior in such a way that revictimization is facilitated: The traumatogenic model (Finkelhor and Browne, 1985); The posttraumatic stress model (Briere, 1997); Learning theory (Messman and Long, 1996, cited in Grauerholz, 2000).

According to Grauerholz, all these frameworks share a basic assumption that CSA can profoundly alter an individual's personality and social development. Research has shown that CSA

victims exhibit traits of traumatic sexualization, such as earlier onset of sexual relations, greater numbers of partners, and prostitution. Other factors include drug and alcohol abuse, low self-esteem, powerlessness, and stigmatization. Grauerholz notes that the latter three factors have not been tested empirically.

Grauerholz also describes the “learned expectancy” idea of Messman and Long (1996). It is suggested that women who have experienced sexual abuse as children “learn and adopt an inappropriate repertoire of sexual behavior, and may learn to associate sexuality with pain, punishment, or other negative consequences” (Messman and Long, 1996, p. 399, cited in Grauerholz, 2000). Thus, such victims may learn that a degree of pain, coercion, and trauma is actually a normal aspect of sexual relations. Such a perspective would certainly give CSA victims a “higher threshold of tolerance” for coercion, force, intimidation, pleading, and lack of affection/regard in their later sexual relations.

The *microsystem* is the immediate (interpersonal) context in which revictimization occurs. Two ways a victim of CSA may become vulnerable to revictimization later in adulthood are suggested. “First, certain factors may lead to increased contact with potential perpetrators, contributing to an exposure risk. Second, certain factors may increase the likelihood that potential perpetrators with whom they come into contact will act aggressively” (Grauerholz, 2000, p. 10). Self-esteem, too, may play a role. Some CSA victims experience low self-esteem, which may lead to greater intimacies with potential abusers “in an effort to verify their self-image” (p. 10).

Aside from the victim, there may be factors that increase the likelihood of perpetrators acting aggressively; signals that lead them to believe their advances will be successful. For example, social isolation, low self-esteem, lack of family support systems, timidity, reluctance, or inability to respond assertively to unwanted sexual advances.

The *Exosystem* level refers to those social systems in which the person is embedded (such as school, church, neighborhood, and extended family). Grauerholz invokes the concept of social power to suggest that CSA victims (who may become

socioeconomically disadvantaged because of early pregnancy, less education, single parenthood, and poverty) are relatively powerless at the exosystem level. Social isolation and poverty, for instance, may shape the opportunities a young female single parent perceives herself as having in terms of dating partners. She may also be less powerful in her intimate relationships and so be more vulnerable to violence from partners.

At the final, *macrosystem*, level Grauerholz considers the larger cultural context in which revictimization occurs. She suggests societal tendencies to “blame the victim” of sexual assaults may facilitate later revictimization.

So, in plain terms, a proportion of victims of sexual abuse (not all) carry profound psychological distress into adulthood. This sadness, distrust, lack of confidence, and low self-esteem makes a certain number vulnerable to revictimization and high-risk sexual behavior, including forms of commercial sexual exploitation, such as prostitution. A vulnerability to substance abuse enhances their risk of being assaulted or becoming involved in sex work. The experience of CSA appears to remove barriers to involvement in sex work for some victims.

CSA: Prevention and Therapy

This section will review the literature on both prevention and therapy in the CSA field.

Prevention

The literature on the prevention of sexual abuse advocates an ecological holistic approach that aims to address the various systemic levels that influence the child’s development (Browne, Hanks, Stratton, & Hamilton-Giachritsis, 2002; Save the Children Norway, 2005; WHO, 2002, 2004, 2006, 2007).

A proper prevention policy needs to include cooperation between the educational, health, social, child protection, police, and judicial sectors and target not only children at risk but also the potential offender. A successful strategy has to attack the problem on different fronts in a holistic manner (Save the Children Norway, 2005, p. 85).

Save the Children Norway (2005) describes several multi-sectoral initiatives from different continents and suggests that most governments have failed to take a central role in coordinating and implementing policies and services for prevention and child protection. WHO (2006) provides guidelines for implementing outcome evaluations of prevention programs of child maltreatment. However, most efforts are directed at one level alone, for example, media campaigns that target the general public, school programs that target children (and to a lesser extent parents), and therapeutic services.

Media Campaigns. Hoefnagels and Baartman (1997) and Hoefnagels and Mudde (2000) report on a multimedia campaign in the Netherlands in 1991–1992, which resulted in increased disclosure of sexual abuse as measured by calls to a children’s

helpline before and after the campaign. Chalk and King (1998) evaluated a CSA media campaign using interviews with 200 parents from 8 U.S. sites and found it effective in terms of increased knowledge and having a positive impact on how people would respond to children reporting sexual abuse. No differences in attitudes toward victims or offenders were noted. The need for raising public awareness of sexual abuse is underscored by such studies as Chen, Dunne, and Han (2007) who found significant proportions of parents who did not know that children most often are sexually abused by familiar people, that females can be the perpetrator, that boys can be sexually abused, and that usually there are no obvious physical signs when sexual abuse has occurred. Rheingold and colleagues (2007), however, stress that media campaigns alone may not significantly affect primary prevention of CSA.

School-Based Prevention Programs. Studies of school-based CSA prevention programs have repeatedly found improvements in the acquisition of knowledge and in some cases, evidence of safety skills, although as many authors note, these improvements do not necessarily reduce the occurrence of sexual abuse. The meta-analysis of Rispens, Aleman, and Goudena (1997) also supported this finding, particularly in relation to preschool children but noted that knowledge gained decreased over time.

Zwi et al. (2007) conducted a Cochrane Review of 15 randomized controlled trials of school-based prevention programs, predominantly conducted in the United States. Although the review of Zwi et al. found that the majority of studies reported significant improvements in knowledge measures and protective behaviors in simulated at-risk situations, only a few studies aimed to change actual behavior, although these studies did find changes in demonstrating safe behaviors. Zwi et al. advocate caution in interpreting the results from their review. They were concerned about both the methodologies used and the analysis involved in the original studies, and the lack of follow-up in terms of whether information was retained over time. Parents have expressed concern about the potential harm of such programs, such as children knowing too much about sex (Chen et al., 2007) that may act as a potential barrier to introducing prevention programs in schools (MacIntyre & Carr, 1999). Surprisingly, few studies appear to have addressed parents’ perceptions of school-based sexual abuse prevention programs.

None of these reviews provide evidence that school-based programs reduce the incidence of sexual abuse or lead to easier access to services for children who have been abused. Outstanding issues highlighted by Zwi et al. (2007) requiring further exploration are the optimal age for interventions, whether knowledge gained is retained in the longer term, whether knowledge and skills learned transfer to real-life situations, and the optimal duration of such programs. A criticism of such programs by parents is that they inadequately address intrafamilial sexual abuse (McElvaney, 2008) and place an inappropriate burden on the child (Save the Children Norway, 2005). Teachers’ attitudes to implementing such programs are important, given the significant variation in teacher comfort in dealing with sexuality issues in the classroom (Mayock, Kitching, & Morgan, 2007).

The scientific literature is conclusive that school prevention programs are effective in dealing with other social issues such as preventing bullying and drug use (see Durlak, 1995, cited in Finkelhor, Ormrod, & Turner, 2007). However, one key distinction between CSA and bullying or drug use is that in sexual abuse, the child is most often the victim of such behavior, rather than the perpetrator. Behavior change is therefore an explicit goal of prevention programs aimed at preventing bullying and drug use and is often measured in evaluations of such programs. The consensus in the sexual abuse literature is that although school prevention programs have been found to be effective in increasing children's knowledge and skills, there is no evidence from outcome evaluation studies to suggest that these skills are transferred to real-life situations and therefore prevent sexual abuse from occurring in the first place. Such outcome measures would be difficult to obtain. Nevertheless, epidemiological studies are reflecting a decline in rates of substantiated CSA in the United States, Canada, and the United Kingdom (Finkelhor, 2008) and school-based prevention programs have been cited as a possible explanation for this decline (Finkelhor & Jones, 2006; Jones, Finkelhor, & Halter, 2006).

Prevention Through Therapy With Abusers. WHO (2006) note the value of therapeutic intervention with various groups as part of a multipronged approach to prevention of child abuse. Secondary prevention is noted as the objective of many therapeutic programs for offenders. Cognitive-behavior therapy (CBT) has been found to be effective, particularly those approaches that include relapse prevention. Hall (1995, cited in Craig, Browne, & Stringer, 2003) found a small but robust treatment effect in his meta-analysis. Alexander (1999, cited in Saunders, Berliner, & Hanson, 2004) reviewed 79 treatment studies and noted recidivism rates of 8.1% compared to 18.3% for other treatment approaches and 25.8% for untreated molesters.

Another area of prevention that has received attention in recent decades is that of targeting adolescent offenders to prevent these adolescents abusing in adulthood, given that a large proportion of sex offenders are adolescents (Synder, 2000; Snyder & Sickmund, 2006, cited in Finkelhor et al., 2007). Outcome studies are promising, highlighting the potential of therapeutic intervention with adolescents as a way of preventing sexual abuse (O'Reilly, Carr, Marshall, & Beckett, 2004).

Key Components of an Effective Preventive Strategy. Save the Children Norway (2005) recommends that governments take a leading role in coordinating a child rights-based national policy framework to address CSA with enforceable agreements between governmental and nongovernmental agencies. According to this report, such a framework should include a strong monitoring and evaluation component, a multi-sectoral and multidisciplinary integrated response, an emphasis on children's consultation processes, a centralized data-gathering and research service, and integrated services for children, their families, and young and adult offenders, which provide immediate response in child protection, multilevel care and, finally,

sufficient resources allocated to such measures. Butchart and colleagues (WHO, 2004) have described strategies for preventing child maltreatment taking account of various developmental stages and levels of intervention at the individual, relationship, societal, and community levels. An example of such a multilevel multi-sectoral approach to preventing child maltreatment in general (as opposed to CSA in particular) is that of the Triple P Positive Parenting Program (Sanders, Markie-Dadds, & Turner, 2003, cited in WHO, 2006). This program, developed in Australia, involves provision of media messages on positive parenting and information resources such as advice sheets and videos, short, targeted intervention for children with specific behavioral problems offered by primary care practitioners, an intensive training program for parents, and services to address broader family issues such as relationship conflict, parental depression, anger, and stress. A number of independent outcome evaluations have shown it to improve family management techniques, parental confidence in effective child rearing, and behavioral outcomes including behavior and aggression in many parts of Europe, including Germany, Switzerland and United Kingdom, China, Hong Kong, New Zealand, and the United States.

A difficulty with the literature on broad-based child maltreatment prevention studies is discerning the impact of such programs on the prevention of sexual abuse as those risk factors specific to sexual abuse may be neglected in such programs. However, these programs may have the potential for adaptation to specifically target the prevention of CSA. The literature is not lacking in helpful models and guidelines to inform prevention. It is unfortunately lacking in evidence for implementation of such models and in efforts to robustly evaluate those that have been implemented.

Therapeutic Intervention

Therapeutic intervention is only part of the response needed to children and families when sexual abuse has occurred. How child protection systems and legal institutions need to respond is discussed in the WHO World Violence Report (WHO, 2002) and guidelines for best practice in investigating allegations of sexual abuse are available in the United States (American Professional Society on the Abuse of Children [APSAC], 1999) and the United Kingdom (Home Office, 2002). The WHO report suggests that evidence is poor in support of therapy with children and families. This is not a fair assessment of the growing body of evidence to support the use of psychotherapy with children who have been sexually abused.

Children who have been sexually abused are a heterogeneous group, often with little in common other than their experience of sexual abuse. Significant proportions of young people do not appear to need therapeutic intervention following an experience of sexual abuse (Edmond, Auslander, Elze, & Bowland, 2006; Saywitz, Mannarino, Berliner, & Cohen, 2000). Research on children who have been sexually abused is typically based on clinical and legal samples that are highly selective, based on therapeutic need and in the case of legal samples, those

willing to pursue for the most part a lengthy and potentially distressing process (McElvaney, 2008). In addition, Goodman-Brown, Edelstein, Goodman, Jones, & Gordon (2003) found that families were less likely to agree to participate in research if the abuse experienced by the child was intrafamilial. There is some evidence to suggest that children abused by someone within the family represent a distinctly different group from those abused outside the family. Such features as an increased reluctance to disclose abuse (DiPietro, Runyan, & Fredrickson, 1997; Hershkowitz, Horowitz, & Lamb, 2005), and a higher likelihood of recantation (Malloy, Lyon, & Quas, 2007), may militate against inclusion in therapy outcome studies. In addition, there is some evidence to suggest that those who have abused within the family show less improvement following therapy (Hetzel-Riggin, Brausch, & Montgomery, 2007) and may be more subject to the cumulative impact of polyvictimization (Finkelhor, et al., 2007) due to exposure to both sexual and emotional abuse. Psychotherapy outcome studies for the most part involve samples of children who have been sexually abused, without distinguishing between those abused within the family or by someone outside the family. Therefore, the review below focuses on effectiveness of therapy with sexual abuse samples, not specifically with intrafamilial abuse samples.

Psychotherapy Effectiveness. Much research on psychotherapy effectiveness with children who have been sexually abused is concerned with comparing different modalities of therapy. Evidence has been found for various therapeutic interventions in review studies ranging from CBT and abuse-specific therapy (Saunders et al., 2004; Saywitz et al., 2000), art therapy (Kolko, 1987), family therapy (Silovsky & Hembree-Kigin, 1994), group therapy (Silovsky & Hembree-Kigin, 1994), and psychodrama (Avinger & Jones, 2007). Trowell et al. (2002) provided evidence for both psychoanalytic psychotherapy and psychoeducational group therapy in a randomized controlled trial study.

Meta-analyses of group treatment outcomes for sexually abused children and adolescents found a significant effect size for group treatments (Hetzel-Riggin et al., 2007; Reeker, Ensing, & Elliott, 1997), as did a meta-analysis of play therapy studies (Bratton, Ray, Rhine, & Jones, 2005).

Hetzel-Riggin et al. (2007) looked at both the effect of treatment modality and also investigated different secondary problems such as behavior problems and psychological distress. They concluded that therapy was more effective than no therapy with play therapy being more effective for social functioning. Cognitive-behavioral, abuse-specific, and supportive therapy in either group or individual formats was the most effective for behavior problems, while cognitive-behavioral, family, and individual therapy (modality unspecified) was most effective for psychological distress. Abuse-specific, cognitive-behavioral, and group therapy was found to be most effective for low self-concept. The authors conclude that the choice of therapy modality should depend on the child's main presenting secondary problem.

Outcomes in general for children who have been sexually abused are more optimistic for those who have parental support (Everson, Hunter, Runyon, Edelsohn, & Coulter, 1989; Trickett, 1997). Chandy, Blum, and Resnick (1997) in their survey of adolescent boys found that one of the most powerful predictors of resilience was the perception that their parents cared about them. Other researchers have found support for concurrent intervention with abused children and their non-abusing parents, in groups or individual sessions, with concurrent parent-child sessions (Cohen, Deblinger, Mannarino, & Steer, 2004; Putnam, 2003; Ramchandani & Jones, 2003; Reeker et al., 1997). Other areas requiring further research are the role of prosocial peer relationships as a potential buffer to the effects of child abuse and neglect (Cicchetti, Toth, & Maughan, 2000) and how understanding contextual characteristics and the mediation of social supports may affect the sequelae of child maltreatment (Pepin & Banyard, 2006).

Evidence From Generic Psychotherapy Literature. The known prevalence of sexual abuse among clinical populations and the underreporting of sexual abuse both underscore the relevance of the generic psychotherapy outcome literature to the field of CSA. This literature provides ample evidence of the effectiveness of psychotherapy as an intervention with children and adolescents with psychological difficulties that may arise from many sources, one of which is frequently CSA. Carr (2007) cites four broad meta-analyses of studies involving children with a broad range of psychological difficulties receiving a variety of different interventions, which included 350 treatment outcome studies. Effect sizes ranged from .71 to .79, indicating that the average child receiving therapy fared better than 76% to 81% of children in control groups (Casey & Berman, 1985; Kazdin, Bass, Ayers, & Rodgers, 1990; Weisz, Weiss, Alicke, & Klotz, 1987; Weisz, Weiss, Han, Granger, & Morton, 1995). Two of these studies found that effect sizes were maintained at 6 months' follow-up (Weisz et al., 1987, 1995). Carr (2007) also found a "dose effect" relationship in psychotherapy, suggesting that 20–45 sessions are necessary for 50–75% of psychotherapy clients to recover.

As with the literature on therapy with children who have been sexually abused, various therapeutic models have been found to be effective with clinical populations. Such findings emanate from reviews of meta-analyses on CBT (Butler, Chapman, Forman, & Beck, 2006) reviews of both controlled studies and those without control groups on psychoanalytic psychotherapy in Germany (Richter et al., 2002; Loew et al., 2002, respectively, cited in Carr, 2007) and in the United Kingdom (Fonagy et al., 2002; Kennedy, 2004) and a review of meta-analyses of family therapy studies (Shadish & Baldwin, 2003, cited in Carr, 2007). These findings support the use of psychotherapy as an empirically based intervention with children and adolescents with psychological difficulties.

The ecological approach advocated in the area of prevention has equally valid application to the therapeutic response to sexual abuse but is rarely used to inform intervention. Levendosky and Buitenheim (2000) report on a single case study where the

therapy was based on an integrated relational and trauma theory perspective through a systemic approach working directly within the family, school, and social service agencies. Although examples of working with both the victim and the offender are not uncommon in clinical practice, the mutual benefits of such work to both victim and offender, particularly in cases of intrafamilial sexual abuse, have received little attention in the research literature.

Hetzel-Riggin et al. (2007) note that because of the heterogeneity of samples in their review, this logically suggests that some therapy modalities may be more effective than others depending on the presenting secondary problems. Saunders et al. (2004) note that the interventions with the most empirical support tend to be based on behavioral or cognitive behavioral theoretical approaches, use behavioral and cognitive intervention procedures and techniques, and intervene at both the individual child and the parent/family levels. Most of the programs reviewed are based on a cognitive behavioral approach with little reference to other modalities. The current thinking in psychotherapy is that common factors across psychotherapy modalities have a greater impact on whether clients benefit from psychotherapy than specific factors (Carr, 2007). According to Carr (2007), the most important single common factor in psychotherapy outcome is the therapeutic relationship or working alliance, where the therapist is empathic and collaborative and the client is cooperative and committed to recovery. Carr (2007) notes that one of the challenges of examining the evidence for psychotherapy is the caution required in concluding that because CBT has been more frequently evaluated than other interventions, it is therefore more effective. Macdonald, Higgins, and Ramchandani (2006) conducted a Cochrane Review of the evidence base for cognitive behavioral interventions with children and adolescents who have been sexually abused. Their review, involving 10 randomized control studies, while confirming the potential of CBT (particularly trauma-focused approaches) as a means of addressing the adverse consequences of CSA, highlighted the tenuousness of the evidence base and the need for more carefully conducted and better reported trials.

The Child's Perspective. Recent developments in the field of sexual abuse research have emphasized the importance of involving children directly in the research process (McElvaney, 2008). Mudaly and Goddard (2006) asked children directly about their experiences of a sexual abuse intervention service, which included both assessment and therapeutic services. Young people's views about child protection and police intervention highlight the complexity of responding to young people's needs in ways that do not render them even more helpless following experiences of abuse. The use of qualitative methodologies can help identify themes or areas of investigation of importance to children themselves.

Reviews of the literature have pointed to the methodological difficulties associated with research studies—small samples with corresponding low predictive power, lack of detail in relation to study design and content of therapeutic intervention, and

lack of control group. As with the ecological approach advocated in the prevention literature, a centrally coordinated, well-resourced holistic approach to therapeutic outcome evaluation is needed to ensure that limited resources are allocated to meeting the needs of children and families who have been affected by CSA. Saunders et al. (2004) offer 22 general principles of treatment for children, which emphasize the importance of conducting comprehensive needs-based assessments, providing evidence-based interventions, including parents and caregivers in the therapeutic work, offering support to other family members where indicated, including the offender and matching the needs of clients to the intervention offered.

Cultural Issues and Accessing Help. Very little information is available on whether cultural differences exist in the relative efficacy of different therapeutic interventions (Cohen, Deblinger, Mannarino, & de Arellano, 2001). Cohen et al. 2001 do suggest that certain cultural groups may have value systems that encourage them to engage with psychological services and influence expectations of therapy. Families from all cultural backgrounds may need help in accessing therapy services. In one U.S. study conducted through telephone interviews with 157 parents whose children had suffered a serious sexual or physical assault in the previous year, only 22% of these parents had even considered getting mental health treatment for their victimized children (Kopiec, Finkelhor, & Wolak, 2004). According to Carr's (2007) extensive review, clients who are able to access therapy more quickly (and who spend little time on waiting lists) are more likely to engage in therapy. Some authors have suggested that cultural differences and associated attitudes toward purity/virginity and/or stigmatization of those sexually victimized influence whether children are more or less likely to disclose experiences of sexual abuse (Kazarian & Kazarian, 1998; Muntarhorn, 1990; Toukmanian & Brouwers, 1998). According to Saunders et al. (2004), the role of cultural, racial, ethnic, and religious identification on treatment response in cases of child abuse is an important area for future research and needs to be considered in all treatment planning decisions.

Conclusion

Notwithstanding significant methodological limitations, the extant literature reveals a growing body of well-designed studies from across the globe indicating the need to view CSA as an international public health problem. The Stockholm (1996) and Yokohama (2001) World Congress against Commercial Sexual Exploitation of Children (CSEC) focused on commercial forms of child sexual exploitation, such as sex tourism, prostitution, pornography, and trafficking. There is a recognition that the much more pervasive sexual abuse of children in their own families and communities, by perpetrators trusted and known to them, needs to be addressed as part of the global initiative to tackle CSA. For example, a 1996 Eastern and Southern African Regional Consultation on CSEC observed that, apart from commercial sexual abuse of children,

there were more alarming occurrences of the noncommercial sexual abuse of children in the form of domestic violence and incest . . . far more attention needs to be paid to abuse in the home as both a primary cause of commercial sexual exploitation and as a far more serious problem in itself. (UNICEF & ANNPCAN, 2001, p. 8)

The identity of perpetrators is of considerable importance. Victims are far less likely to report CSA when it is perpetrated by somebody close to them for reasons of fear of perpetrator, fear of consequences, embarrassment, and shame (see Pipe, Lamb, Orbach, & Cederborg, 2007 for a review on inhibitors to disclosure). This increases the likelihood of environmental systems, such as police, hospitals and courts becoming aware of abuse by strangers to a disproportionate degree.

Evidence that the sexual abuse of children is universally widespread, and is most frequently perpetrated by family members, neighbors, peers, and others known to the child or adolescent is indisputable. CDCP and UNICEF (2007) provide a strong argument for the need to have good data about CSA:

In the absence of good prevalence estimates the tendency in many countries is to believe that childhood sexual violence is not a problem. Further, progress towards prevention cannot be monitored without reliable population based baseline and follow-up data. Population based data can also be used to identify potential risk and protective factors for sexual violence in order to develop effective prevention strategies. (p. 11)

The short- and long-term effects of CSA are variable, depending on the age of onset, number and identity of perpetrators, severity, degree of family support, and other factors. In addition to socioemotional and mental health sequelae, studies have consistently found that victims are vulnerable to subsequent sexual revictimization in adolescence and adulthood. Russell's 1986 study suggested that the risk is twofold for women who experienced intrafamilial sexual abuse before the age of 14 for rape or attempted rape after the age of 14, while a review of 19 empirical studies of Roodman and Clum (2001) found an effect size of .59 when including broader definitions of CSA.

Studies have also found a link between CSA and later engagement in high-risk sexual behavior, prostitution and sexual promiscuity. Individuals who have experienced CSA are also more likely than non-victims to engage in potentially harmful and damaging high-risk sexual behaviors, such as early age at first intercourse, multiple partners, unprotected sex resulting in unplanned pregnancies or sexually transmitted diseases, and prostitution. Although this relationship cannot be considered a causal relationship, there is an abundance of evidence that the experience of childhood sexual abuse, in particular penetrative abuse, significantly increases the risk for such high-risk sexual behaviors. More severe forms of sexual abuse and the coexistence of physical abuse increases the risks for later revictimization and engagement in high-risk sexual behaviors.

Mediating factors such as parental psychopathology, family conflict versus a supportive family environment, and leaving

home at an early age appear to play a role in these relationships. However, there is a need for research that separates out the confounding influence of such variables to realize the true influence of such mediators.

Cross-sectional studies cannot explain these outcomes. Although theoretical explanatory models that focus on the psychosocial impacts of the abuse on the individual have been commonly relied on to enhance our understanding of these links, these models neglect contextual factors which according to some authors are equally important in considering such relationships. Ecological models, such as that proposed by Grauerholz (2000), that incorporate explanations at the individual, family, community, and societal level, are promising. Grauerholz emphasizes the reciprocal nature of influences whereby individual factors such as psychological coping strategies interact with environmental factors such as exposure to family environments and high-risk situations leads to the greater risk of revictimization and engagement in high-risk sexual behaviors. Grauerholz' theory is consistent with a changing focus in the literature from the individual to interpersonal and cultural factors within society.

Interventions in the area of prevention of CSA have not kept pace with international policy guidelines. Thus, whether such guidelines have empirical support is open to question. Nevertheless, preliminary attempts at mass media campaigns and school-based prevention programs have shown some improvements in the increase of knowledge and to a more limited degree, the acquisition of skills. As yet, there is no evidence that interventions have prevented sexual abuse from occurring in the first place.

However, broad-based prevention programs aimed at preventing child maltreatment in general (as opposed to CSA in particular) may have something to offer in informing CSA prevention programs. Centralized responsibility for coordinating such a broad-based approach appears to be crucial to its success. Although an ecological approach is promoted in the area of prevention, the current literature review highlights the need for such an approach in the area of therapy—with the child who has been abused, the family (which may or may not include the abuser), and the community. To date, such an approach has not been evident in practice or in the research literature. There is considerable evidence to support the use of various therapeutic modalities in intervening with children and families following the experience of CSA. However, researchers have identified significant methodological limitations in the extant research literature that impede the making of recommendations for implementing existing therapeutic programs unreservedly.

We conclude by quoting CDCP and UNICEF (2007), who state “[sexual] violence against children is both a public health problem and a human rights problem” (p. 28). The former because of the negative sequelae experienced in the population of victims (far too high a population in many societies) and the latter because of the infringement of the personal integrity of the victim.

Note

1. The phenomenon of “transactional sex” must be considered in the African context. Sexual relations between younger female adolescents and older boys or men, with the exchange of small gifts or money, are frequently exploitative. García-Moreno and Watts (2000) refer to this as “economically coerced sex” (p. 261). Older boys/men who provide financial/economic support to young girls are widely known as “sugar daddies.” Although most likely to occur in conditions of poverty and deprivation, such relationships have also been noted among the middle and privileged classes, where additional funds may be used for fashionable clothes, mobile phones, and other luxury items. Silberschmidt (2001) argues that the patterns of sexual behavior evident in parts of sub-Saharan Africa today are not the result of some traditional “permissiveness,” but the result of the breakdown of traditional norms and regulations surrounding sexual behavior, aggravated by widespread poverty and social upheaval.

Acknowledgments

This article is based on a review commissioned by ISPCAN for the Pre “3rd World Congress on Sexual Exploitation” Workshop (Hong Kong, September 2008). Resources can be found at www.ispcan.org/congress2008/wcse.html. The authors are grateful for the useful comments from reviewers and to Dublin Institute of Technology for its support.

Declaration of Conflicting Interests

The author(s) declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

This review was funded by a grant from UNICEF (Geneva).

References

- American Professional Society on the Abuse of Children. (1999). *Guidelines for psychosocial evaluation of suspected sexual abuse in young children*. Chicago, IL: Author.
- Arata, C. M. (2000). From child to adult victim: A model for predicting sexual revictimization. *Child Maltreatment, 5*, 1, 28-38.
- Arata, C. M. (2002). Child sexual abuse and sexual revictimization. *Clinical Psychology: Science and Practice, 9*, 135-164.
- Avinger, K. A., & Jones, D. H. (2007). Group treatment of sexually abused adolescent girls: A review of outcome studies. *American Journal of Family Therapy, 35*, 315-326.
- Bagley, C., & Young, L. (1987). Juvenile prostitution and child sexual abuse: A controlled study. *Canadian Journal of Community Mental Health, 6*, 5-26.
- Belsky, J. (1980). Child maltreatment: An ecological integration. *American Psychologist, 35*, 320-335.
- Bratton, S. C., Ray, D., Rhine, T., & Jones, L. (2005). The efficacy of play therapy with children: A meta-analytic review of treatment outcomes. *Professional Psychology: Research and Practice, 36*, 376-390.
- Briere, J. (1992). Methodological issues in the study of sexual abuse effects. *Journal of Consulting and Clinical Psychology, 60*, 196-203.
- Briere, J. (1997). *Psychological assessment of adult posttraumatic states*. Washington, DC: American Psychological Association.
- Briere, J., & Elliott, D. (2003). Prevalence and psychological sequelae of self-reported childhood physical and sexual abuse in a general population sample of men and women. *Child Abuse and Neglect, 27*, 1205-1222.
- Briere, J. & Runtz, M. (1993). Childhood sexual abuse: Long-term sequelae and implications for psychological assessment. *Journal of Interpersonal Violence, 8*, 312-330.
- Bronfenbrenner, U. (1979). *The ecology of human development*. Cambridge, MA: Harvard University Press.
- Browne, K. D., Hanks, H., Stratton, P., & Hamilton-Giachritsis, C. (Eds.). (2002). *Early prediction and prevention of child abuse: A handbook*. Chichester, UK: John Wiley.
- Butler, A., Chapman, J., Forman, E., & Beck, A. (2006). The empirical status of cognitive-behavioural therapy: A review of meta-analyses. *Clinical Psychology Review, 26*, 17-31.
- Carr, A. (2007). *The effectiveness of psychotherapy: A review of research*. Dublin: Irish Council of Psychotherapy.
- Casey, R. J., & Berman, J. S. (1985). The outcome of psychotherapy with children. *Psychological Bulletin, 98*, 388-400.
- CDCP & UNICEF. (2007). *Violence against children in Swaziland: Findings from a National Survey on Violence Against Children in Swaziland*. Author, location: Mbabana.
- Chalk, R., & King, P. (1998). Assessing family violence interventions. *American Journal of Preventive Medicine, 14*, 289-292.
- Chandy, J. M., Blum, R. W., & Resnick, M. D. (1997). Sexually abused male adolescents: How vulnerable are they? *Journal of Child Sexual Abuse, 6*, 1-16.
- Chen, J. Q., Dunne, M., & Han, P. (2004). Child sexual abuse in China: A study of adolescents in four provinces. *Child Abuse & Neglect, 28*, 1171-1186.
- Chen, J. Q., Dunne, M., & Han, P. (2007). Prevention of child sexual abuse in China: Knowledge, attitudes, and communication practices of parents of elementary school children. *Child Abuse & Neglect, 31*, 747-755.
- Cicchetti, D., Toth, S., & Maughan, A. (2006). An ecological-transactional model of child maltreatment. In A. Samaroff, M. Lewis & S. Miller (Eds.), *Handbook of developmental psychopathology*. Dordrecht, the Netherlands: Kluwer Academic Publishers.
- Classen, C., Palesh, O., & Aggarwal, R. (2005). Sexual revictimization: A review of the literature. *Trauma, Violence and Abuse, 6*, 103-129.
- Cohen, J. A., Deblinger, E., Mannarino, A. P., & de Arellano, M. A. (2001). The importance of culture in treating abused and neglected children: An empirical review. *Child Maltreatment, 6*, 148-157.
- Cohen, J. A., Deblinger, E., Mannarino, A. P., & Steer, R. A. (2004). A multisite, randomized controlled trial for children with sexual abuse-related PTSD symptoms. *Journal of the American Academy of Child and Adolescent Psychiatry, 43*, 393-402.
- Collishaw, S., Pickles, A., Messer, J., Rutter, M., Shearer, C., & Maughan, B. (2007). Resilience to adult psychopathology following childhood maltreatment: Evidence from a community sample.

- Child Abuse & Neglect*, 31, 211-229.
- Craig, L. A., Browne, K. D., & Stringer, I. T. (2003). Treatment and sexual offense recidivism. *Trauma, Violence, & Abuse*, 4, 70-89.
- DeMause, L. (1991). The universality of incest. *The Journal of Psychohistory*, 19, 123-164.
- DiPietro, E. K., Runyan, D. K., & Fredrickson, D. D. (1997). Predictors of disclosure during medical evaluation for suspected sexual abuse. *Journal of Child Sexual Abuse*, 1, 133-142.
- Dube, S., Anda, R., Whitfield, C., Brown, D., Felitti, V., Dong, M., & Giles, W. (2005). Long term consequences of childhood sexual abuse by gender of victim. *American Journal of Preventative Medicine*, 28, 430-438.
- Dunkle, K., Jewkes, R., Brown, H., Yoshihama, M., Gray, G., McIntyre, J., & Harlow, S. (2004). Prevalence and patterns of gender based violence and revictimization among woman attending antenatal clinics in Soweto, South Africa. *American Journal of Epidemiology*, 160, 230-239.
- Edmond, T., Auslander, W., Ezle, D., & Bowland, S. (2006). Signs of resilience in sexually abused adolescent girls in the foster care system. *Journal of Child Sexual Abuse*, 15, 1-28.
- Elbedour, S., Abu-Bader, S., Onwuegbuzie, A., Abu-Rabia, A., & El-Aassam, S. (2006). The scope of sexual, physical, and psychological abuse in a Bedouin-Arab community of female adolescents: The interplay of racism, urbanization, polygamy, family honor, and the social marginalization of women. *Child Abuse & Neglect*, 30, 215-229.
- Everson, M. D., Hunter, W. M., Runyon, D. K., Edelsohn, G. A., & Coulter, M. L. (1989). Maternal support following disclosure of incest. *American Journal of Orthopsychiatry*, 59, 197-207.
- Fanslow, J. L., Robinson, E., Crengle, S., & Perese, L. (2007). Prevalence of child sexual abuse reported by a cross-sectional sample of New Zealand women. *Child Abuse & Neglect*, 31, 935-945.
- Fergusson, D., Horwood, J., & Lynskey, M. (1997). Childhood sexual abuse, adolescent sexual behaviours and sexual revictimization. *Child Abuse and Neglect*, 21, 789-803.
- Finkelhor, D. (1990). Early and long term effects of child sexual abuse: An update. *Professional Psychology: Research and Practice*, 21, 325-330.
- Finkelhor, D. (1994). The international epidemiology of child sexual abuse. *Child Abuse & Neglect*, 18, 409-417.
- Finkelhor, D. (2008). *Lessons from the decline in sexual abuse*. Paper presented at the St. Clare's Unit & St. Louise's Unit Joint Annual Conference, Dublin.
- Finkelhor, D., & Browne, A. (1985). The traumatic impact of child sexual abuse. *American Journal of Orthopsychiatry*, 55, 530-541.
- Finkelhor, D., & Jones, L. (2006). Why have child maltreatment and child victimization declined? *Journal of Social Issues*, 62, 685-716.
- Finkelhor, D., Ormrod, R. K., & Turner, H. A. (2007). Polyvictimization: A neglected component in child victimization. *Child Abuse & Neglect*, 31, 7-26.
- Fleming, J., Mullen, P., Sibthorpe, B., & Bammer, G. (1999). The long term impact of childhood sexual abuse in Australian women. *Child Abuse & Neglect*, 23, 145-159.
- Flowers, R. (1998). *The prostitution of women and girls*. Jefferson, NC: McFarland.
- Fonagy, P., Clarkin, J., Gerber, A., Kachele, H., Krause, R., & Jones, E. E. (2002). *Open door review of outcome studies in psychoanalysis*. London: International Psychoanalytic Association.
- García-Moreno, C., & Watts, C. (2000). Violence against women: Its importance for HIV/AIDS. *AIDS*, 14, 253-265.
- García-Moreno, C., Jansen, H., Ellsberg, M., Heise, L., & Watts, C. (2005). *WHO multi-country study on women's health and domestic violence against women*. Geneva: World Health Organization.
- Goodman-Brown, T. B., Edelstein, R. S., Goodman, G. S., Jones, D. P. H., & Gordon, D. S. (2003). Why children tell: A model of children's disclosure of sexual abuse. *Child Abuse & Neglect*, 27, 525-540.
- Grauerholz, L. (2000). An ecological approach to understanding sexual revictimization: Linking personal, interpersonal, and sociocultural factors and processes. *Child Maltreatment*, 5, 5-17.
- Hershkowitz, I., Horowitz, D., & Lamb, M. E. (2005). Trends in children's disclosure of abuse in Israel: A national study. *Child Abuse & Neglect*, 29, 1203-1214.
- Hetzel-Riggin, M. D., Brausch, A. M., & Montgomery, B. S. (2007). A meta-analytic investigation of therapy modality outcomes for sexually abused children and adolescents: An exploratory study. *Child Abuse & Neglect*, 31, 125-141.
- Hoefnagels, C., & Baartman, H. (1997). On the threshold of disclosure: The effects of a mass media field experiment. *Child Abuse & Neglect*, 21, 557-573.
- Hoefnagels, C., & Mudde, A. (2000). Mass media and disclosures of child abuse in the perspective of secondary prevention: Putting ideas into practice. *Child Abuse & Neglect*, 24, 1091-1101.
- Home Office. (2002). *Achieving best evidence in criminal proceedings: Guidance for vulnerable or intimidated witnesses, including children*. London: Her Majesty's Stationery Office.
- Humphrey, J. A., & White, J. W. (2000). Women's vulnerability to sexual assault from adolescence to young adulthood. *Journal of Adolescent Health*, 27, 419-424.
- Jewkes, R., Levin, L., Mbananga, N., & Bradshaw, D. (2002). Rape of girls in South Africa. *The Lancet*, 359, 319-320.
- Jones, L., Finkelhor, D., & Halter, S. (2006). Child maltreatment trends in the 1990s: Why does neglect differ from sexual and physical abuse? *Child Maltreatment*, 11, 107-120.
- Kazarian, S. S., & Kazarian, L. Z. (1998). Cultural aspects of family violence. In S. S. Kazarian & D. R. Evan (Eds.), *Cultural clinical psychology: Theory, research and practice* (pp. 316-347). Oxford: Oxford University Press.
- Kazdin, A., Bass, D., Ayers, W. A., & Rodgers, A. (1990). Empirical and clinical focus of child and adolescent psychotherapy research. *Journal of Consulting & Clinical Psychology*, 58, 729-740.
- Kempe, C., Silverman, F., Steele, B., Droegmueller, W., & Silver, H. (1962). The battered child syndrome. *Journal of the American Medical Association*, 181, 17-24.
- Kendall-Tackett, K. A., Williams, L. M., & Finkelhor, D. (1993). Impact of sexual abuse on children: A review and synthesis of the literature. *Psychological Bulletin*, 113, 164-180.
- Kennedy, E. (2004). *Child and adolescent psychotherapy: A systematic review of psychoanalytic approaches*. London: North Central London, NHS Strategic Health Authority.

- King, G., Guilbert, P., Ward, D., Arwidson, P., & Noubary, F. (2006). Correlates of sexual abuse and smoking among French adults. *Child Abuse & Neglect, 30*, 709-723.
- Kolko, D. J. (1987). Treatment of child sexual abuse: Programs, progress, and prospects. *Journal of Family Violence, 2*, 303-318.
- Kopiec, K., Finkelhor, D., & Wolak, J. (2004). Which juvenile crime victims get mental health treatment? *Child Abuse & Neglect, 28*, 45-59.
- Korbin, J. E. (Ed.). (1981). *Child abuse and neglect: Cross cultural perspectives*. Berkeley: University of California Press.
- Lachman, P. (1996). Child protection in Africa: The road ahead. *Child Abuse & Neglect, 20*, 543-547.
- Lalor, K. (2008). Child sexual abuse and HIV transmission in sub-Saharan Africa. *Child Abuse Review, 17*, 94-107.
- Lalor, K., de Róiste, A. & Devlin, M. (2007). *Young people in contemporary Ireland*. Dublin: Gill and Macmillan.
- Levendosky, A. A., & Buttenheim, M. (2000). A multi-method treatment for child survivors of sexual abuse: An intervention informed by relational and trauma theories. *Journal of Child Sexual Abuse, 9*, 1-19.
- Macdonald, G. M., Higgins, J. P. T., & Ramchandani, P. (2006). Cognitive-behavioural interventions for children who have been sexually abused. *Cochrane Database of Systematic Reviews, 4*, CD001930. doi:10.1002/14651858.CD001930.pub2
- MacIntyre, D., & Carr, A. (1999). Evaluation of the effectiveness of the Stay Safe primary prevention programme for child sexual abuse. *Child Abuse & Neglect, 23*, 1307-1325.
- Malloy, L. C., Lyon, T. D., & Quas, J. (2007). Filial dependency and recantation of child sexual abuse allegations. *Journal of the American Academy of Child and Adolescent Psychiatry, 46*, 162-170.
- Mandoki, C., & Burkhart, B. (1989). Sexual victimization: Is there a vicious cycle? *Violence and Victims, 4*, 179-190.
- Masson, J. (1992). *The assault on truth: Freud and child sexual abuse* (3rd ed.). London: Fontana.
- May-Chahal, C., & Cawson, P. (2005). Measuring child maltreatment in the United Kingdom: A study of the prevalence of child abuse and neglect. *Child Abuse & Neglect, 29*, 969-984.
- Mayock, P., Kitching, K., & Morgan, M. (2007). *RSE in the context of SPHE: An assessment of the challenges to full implementation of the programme in post-primary schools*. Dublin: Crisis Pregnancy Agency/Dept of Education and Science.
- McCann, D., Lalor, K., & Katabaro, K. (2006). Childhood sexual abuse among university students in Tanzania. *Child Abuse & Neglect, 30*, 1343-1351.
- McElvaney, R. (2008). *How children tell: Containing the secret of child sexual abuse* (Unpublished PhD dissertation). Trinity College Dublin.
- McGee, H., Garavan, R., de Barra, M., Byrne, J., & Conroy, R. (2002). *The SAVI report: Sexual abuse and violence in Ireland – A national study of Irish experiences, beliefs and attitudes concerning sexual violence*. Dublin, Ireland: The Liffey Press & Dublin Rape Crisis Centre.
- Messman-Moore, T., & Long, P. (2003). The role of childhood sexual abuse sequelae in the sexual revictimization of women: An empirical review and theoretical reformulation. *Clinical Psychology Review, 23*, 537-571.
- Mudaly, N., & Goddard, C. (2006). *The truth is longer than a lie: Children's experiences of abuse and professional interventions*. London, England: Jessica Kingsley Publishers.
- Muntarhorn, V. (1990). *Sexual exploitation of children*. New York, NY: United National Publications.
- Noll, J., Trickett, P., & Putnam, F. (2003). A prospective investigation of the impact of childhood sexual abuse on the development of sexuality. *Journal of Consulting and Clinical Psychology, 71*, 575-586.
- O'Reilly, G., Carr, A., Marshall, W. L., & Beckett, R. C. (2004). *Handbook of clinical interventions with juvenile sex offenders*. London, England: Taylor and Francis.
- Paine, M. L., & Hansen, D. J. (2002). Factors influencing children to self-disclose sexual abuse. *Clinical Psychology Review, 22*, 271-295.
- Pepin, E. N., & Banyard, V. L. (2006). Social support: A mediator between child maltreatment and developmental outcomes. *Journal of Youth and Adolescence, 35*, 617-630.
- Pereda, N., Guilera, G., Forn, M., & Gomez-Benito, J. (2009). The international epidemiology of child sexual abuse: A continuation of Finkelhor (1994). *Child Abuse and Neglect, 33*, 331-342.
- Pipe, M. E., Lamb, M., Orbach, Y., & Cederborg, A. C. (Eds.). (2007). *Child sexual abuse: Disclosure, delay and denial*. Mahwah, NJ: Lawrence Erlbaum.
- Polusny, M., & Follette, V. (1995). Long-term correlates of child sexual abuse: Theory and review of the empirical literature. *Applied and Preventative Psychology, 4*, 143-166.
- Potter, K., Martin, J., & Romans, S. (1999). Early developmental experiences of female sex workers: A comparative analysis. *Australian and New Zealand Journal of Psychiatry, 33*, 935-940.
- Putnam, F. (2003). Ten-year research update review: Child sexual abuse. *Journal of the American Academy of Child and Adolescent Psychiatry, 42*, 269-278.
- Ramchandani, P., & Jones, D. P. H. (2003). Treating psychological symptoms in sexually abused children: From research findings to service provision. *British Journal of Psychiatry, 183*, 484-490.
- Reeker, J., Ensing, D., & Elliott, R. (1997). A meta-analytic investigation of group treatment outcomes for sexually abused children. *Child Abuse & Neglect, 21*, 669-680.
- Rheingold, A. A., Campbell, C., Self-Brown, S., de Arellano, M. A., Resnick, H., & Kilpatrick, D. (2007). Prevention of child sexual abuse: Evaluation of a community media campaign. *Child Maltreatment, 12*, 352-363.
- Rispers, J., Aleman, A., & Goudena, P. P. (1997). Prevention of child sexual abuse victimization: A meta-analysis of school programs. *Child Abuse & Neglect, 21*, 975-987.
- Runyan, D. (2000). The ethical, legal and methodological implications of directly asking children about abuse. *Journal of Interpersonal Violence, 15*, 675-681.
- Roodman, A., & Clum, G. (2001). Revictimization rates and method variance: A meta-analysis. *Clinical Psychology Review, 21*, 183-204.
- Russell, D. (1986). *The secret trauma: Incest in the lives of girls and women*. New York, NY: Basic Books.
- Sandberg, D., Matorin, A., & Lynn, S. (1999). Dissociation, posttraumatic symptomatology, and sexual revictimization: A prospective

- examination of mediator and moderator effects. *Journal of Traumatic Stress, 12*, 127-138.
- Sanders, B., & Moore, D. (1999). Childhood maltreatment and date rape. *Journal of Interpersonal Violence, 14*, 115-124.
- Saunders, B. E., Berliner, L., & Hanson, R. F. (2004). *Child physical and sexual abuse: Guidelines for treatment (Revised Report: April 26, 2004)*. Charleston, SC: National Crime Victims Research and Treatment Center.
- Save the Children Norway. (2005). 10 Essential learning points: Listen and speak out against sexual abuse of girls and boys. Global submission by The International Save the Children Alliance, UN study on violence against children.
- Saywitz, K. J., Mannarino, A. P., Berliner, L., & Cohen, J. A. (2000). Treatment for sexually abused children and adolescents. *American Psychologist, 55*, 1040-1049.
- Schultz, L. G. (1982). Child sexual abuse in historical perspective. *Journal of Social Work and Human Sexuality, 1*, 21-35.
- Silberschmidt, M. (2001). Disempowerment of men in rural and urban East Africa: implications for male identity and sexual behavior. *World Development, 29*, 657-671.
- Silbert, M., & Pines, A. (1982). Victimization of street prostitutes. *Victimology, 7*, 122-133.
- Silovsky, J. F., & Hembre-Kigin, T. L. (1994). Family and group treatment for sexually abused children: A review. *Journal of Child Sexual Abuse, 3*, 1-20.
- Slonim-Nevo, V., & Mukuka, L. (2007). Child abuse and AIDS-related knowledge, attitudes and behaviour among adolescents in Zambia. *Child Abuse & Neglect, 31*, 143-159.
- Steel, J., & Herlitz, C. (2005). The association between childhood and adolescent sexual abuse and proxies for sexual risk behavior: A random sample of the general population of Sweden. *Child Abuse & Neglect, 29*, 1141-1153.
- Stoltz, J.-A. M., Shannon, K., Kerr, T., Zhang, R., Montaner, J., & Wood, E. (2007). Associations between childhood maltreatment and sex work in a cohort of drug-using youth. *Social Science & Medicine, 65*, 1214-1221.
- Toukmanian, S. G., & Brouwers, M. C. (1998). Cultural aspects of self-disclosure and psychotherapy. In S. S. Kazarian & D. R. Evans (Eds.), *Cultural clinical psychology: Theory, research and practice* (pp. 106-126). New York, NY: Oxford University Press.
- Tricket, P. K. (1997). Sexual and physical abuse and the development of social competence. In S. S. Luthar, J. A. Burack, D. Cicchetti & J. R. Weisz (Eds.), *Developmental psychopathology: Perspectives on adjustment, risk and danger*. New York, NY: Cambridge University Press.
- Trowell, J., Kolvin, I., Weermanthri, T., Sadowski, H., Berelowitz, M., Glaser, D., & Leitch, I. (2002). Psychotherapy for sexually abused girls: Psychopathological outcome findings and patterns of change. *British Journal of Psychiatry, 180*, 234-247.
- UNICEF & ANPPCAN. (2001). *Analysis of the situation of sexual exploitation of children in the Eastern and Southern Africa region*. Retrieved from www.focalpointngo.org/yokohama/latestnews/CSECEastSouthernAfricaDraft
- Weisz, J. R., Weiss, B., Alicke, M. D., & Klotz, M. L. (1987). Effectiveness of psychotherapy with children and adolescents: A meta-analysis for clinicians. *Journal of Consulting & Clinical Psychology, 55*, 542-549.
- Weisz, J. R., Weiss, B., Han, S. S., Granger, D. A., & Morton, T. (1995). Effects of psychotherapy with children and adolescents revisited: A meta-analysis of treatment outcome studies. *Psychological Bulletin, 117*, 450-468.
- Widom, C. (1988). Sampling biases and implications for child abuse research. *American Journal of Orthopsychiatry, 58*, 260-270.
- Widom, C. S., Czaja, S., & Dutton, M. A. (2008). Childhood victimization and lifetime revictimization. *Child Abuse and Neglect, 32*, 785-796.
- Widom, C. S., & Kuhns, J. (1996). Childhood victimization and subsequent risk for promiscuity, prostitution and teenage pregnancy: A prospective study. *American Journal of Public Health, 86*, 1607-1612.
- Wingood, G., & DiClemente, R. (1997). Child sexual abuse, HIV sexual risk, and gender relations of African-American women. *American Journal of Preventative Medicine, 13*, 380-384.
- Worku, D., Gebremariam, A., & Jayalakshmi, S. (2006). Child sexual abuse and its outcomes among high school students in southwest Ethiopia. *Tropical Doctor, 36*, 137-140.
- World Health Organization. (2002). *World report on violence and health*. Geneva, Switzerland: Author.
- World Health Organization. (2004). *Preventing violence: A guide to implementing the recommendations of the World report on violence and health*. Geneva, Switzerland: Author.
- World Health Organization. (2006). *Preventing child maltreatment: A guide to taking action and generating evidence*. Geneva, Switzerland: Author.
- World Health Organization. (2007). *Preventing child maltreatment in Europe: A public health approach*. Geneva, Switzerland: Author.
- Zierler, S., Feingold, L., Laufer, D., Velentgas, P., Kantrowitz-Gordon, I., & Maye, K. (1991). Adult survivors of childhood sexual abuse and subsequent risk of HIV infection. *American Journal of Public Health, 81*, 572-575.
- Zwi, K. J., Woolfenden, S. R., Wheeler, D. M., O'Brien, T. A., Tait, P., & Williams, K. W. (2007). School-based education programmes for the prevention of child sexual abuse (Review). *Cochrane Database of Systematic Reviews, 3*, CD004380.

Bios

Kevin Lalor, PhD, is Head of the Department of Social Sciences, Dublin Institute of Technology. He is the editor of the *Irish Journal of Applied Social Studies* and a member of the editorial board of *Child Abuse and Neglect: The International Journal*. He is secretary to the Royal Irish Academy Social Sciences Committee. Research interests include victimology, child abuse in Africa and youth. He is author (with Áine de Róiste and Maurice Devlin) of *Young People in Contemporary Ireland* (Gill & Macmillan, 2007). In 2008, he presented the International Society for the Prevention of Child Abuse and Neglect (ISPCAN) review article on familial child sexual abuse and

sexual revictimization at the 3rd World Congress against Sexual Exploitation of Children and Adolescents.

Rosaleen McElvaney, PhD, is a Clinical Psychologist and a Lecturer in Developmental/Abnormal Psychology in the Department of Social Sciences at Dublin Institute of Technology. Her clinical work for the

past 20 years has been primarily in the field of child sexual abuse with children, adolescents, and adults. Her doctoral dissertation addressed the issue of how children tell of their experiences of abuse. She was President of the Psychological Society of Ireland in 2008-2009 and sat on the National Board for Safeguarding Children in the Catholic Church in Ireland.